

UNDERSTANDING SCATTERING OF SOUND AT BASIN-SCALES WITH NUMERICAL EXPERIMENTS AND THEORY

John L. Spiesberger
Department of Meteorology
512 Walker Bldg.
University Park, PA 16802

phone: (814) 863-8601 fax: (814) 863-9527 email: jspies@ems.psu.edu

Award #: N00014-95-1-0755

LONG-TERM GOALS

There are two goals of this research. The first is to understand the role of internal waves and the mesoscale in scattering sound in the vertical dimension. The second is to develop computer algorithms to predict the extent of this scattering.

OBJECTIVES

Acoustic transmissions at 133 Hz, 60 ms resolution, were received at 3709 km from the Kaneohe source at a receiver mounted 1000 m beneath the axis of the sound channel. Ray traces and parabolic approximations indicate that the duration of the received energy should be about 2.5 s if the sound propagates through a climatological-average of the sound-speed profile in the northeast Pacific (Spiesberger and Tappert, 1996). But the duration of the acoustic energy is 3.5 s. In order to account for this additional second, we postulate that the meso and fine-scales scatter energy at least 1000 m downwards from the axis of the sound channel.

APPROACH

We utilized the U. Miami Parabolic Equation model to simulate the broadband transmission of sound from the source to the receiver (Tappert *et al.*, 1995). The effects of the mesoscale and internal wave fields on the propagation of sound are estimated separately and together by modelling the sound speed field with climatological, climatological+mesoscale, and climatological+mesoscale+internal wave fields. The research was conducted in collaboration with Dr. Michael Wolfson at Pennsylvania State University.

WORK COMPLETED

Numerical simulations are completed and have been compared with the data (Wolfson and Spiesberger, 1997).

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Understanding Scattering of Sound at Basin-scales with Numerical Experiments and Theory				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Pennsylvania State University, Department of Meteorology, University Park, PA, 16802				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

RESULTS

We found that both the mesoscale and internal wave fields are required to understand the multipath at the receiver. The mesoscale vertically scatters the sound by a few hundred meters and introduces a travel time bias of about 0.6 s for near-axial multipath. The bias delays the arrival times of these multipath. These delayed arrivals are further scattered an additional 800 m down to the depth of the receiver by the internal wave field. The travel time bias introduced by internal waves is relatively small, e.g. $O(10)$ ms.

The observed multipath which are stable at the receiver are also stable in the numerical simulations (Spiesberger and Tappert, 1996; Wolfson and Spiesberger, 1997). The stability of these modelled paths in the presence of fluctuations adds to the body of evidence that these paths are useful for mapping temperature change (Spiesberger *et al.*, 1998).

IMPACT/APPLICATIONS

The fluctuations of eddies and internal waves limit the number of multipath that may be available for mapping the sound speed field with tomography. The near-axial multipath are very sensitive to these fluctuations, and it is not clear how to use them for tomographic inversions. Multipath which are sensitive to fluctuations may be useful for inferring statistical properties of the eddies and internal waves.

TRANSITIONS

No transitions have been identified.

RELATED PROJECTS

The vertical scattering of sound by internal waves has been investigated by Colosi *et al.* (1994). Wolfson (pers. comm.) is investigating the sensitivity of rays to the mesoscale in three-dimensions. M. Brown and F. Tappert (pers. comm.) are investigating ray and finite-wavelength chaos in the presence of ocean structures.

REFERENCES

Colosi, J.A., S. M. Flatte, and C. Bracher, Internal-wave effects on 1000-km oceanic acoustic pulse propagation: Simulation and comparison with experiment, *J. Acoust. Soc. Am.*, 96, 452-468, 1994.

Spiesberger, J.L., Hurlburt, H.E., Johnson, M., Keller, M., Meyers, S., and O'Brien, J.J., Acoustic thermometry data compared with two ocean models: The importance of Rossby waves and ENSO in modifying the ocean interior, in press, *Dynamics of Atmospheres and Oceans*, 26(4), (1998).

Spiesberger, J.L. and Tappert, F., Kaneohe acoustic thermometer further validated with rays over 3700 km and the demise of the idea of axially trapped energy, J. Acoust. Soc. Am., 99, 173-84, (1996).

Tappert, F., J. Spiesberger, and L. Boden, New full-wave approximation for ocean acoustic travel time predictions, J. Acoust. Soc. Am., 97, 2771-2782, 1995.

Wolfson, M. A. and Spiesberger, J. L., Full wave simulation of the forward scattering of sound in a structured ocean: A comparison with observations, submitted to the J. Acoust. Soc. Am., (1997).